



Norbrook[®]
Pharmaceuticals Worldwide



Noromycin 300 LA

Oxytetracycline 300 mg/mL

Noromycin 300 LA is a ready-to-use injectable, broad-spectrum antibiotic. For beef cattle, non-lactating dairy cattle and calves, including veal calves, Noromycin 300 LA is indicated for the treatment of pneumonia and shipping fever complex associated with *Pasteurella* spp., and *Haemophilus* spp., infectious bovine keratoconjunctivitis (pinkeye) caused by *Moraxella bovis*, footrot and diphtheria caused by *Fusobacterium necrophorum*, bacterial enteritis (scours) caused by *E. coli*, wooden tongue caused by *Actinobacillus lignieresii*, leptospirosis caused by *Leptospira pomona*, and wound infections and acute metritis caused by strains of staphylococcal and streptococcal organisms sensitive to oxytetracycline. In swine, Noromycin 300 LA is indicated in the treatment of bacterial enteritis (scours, colibacillosis) caused by *E. coli*, pneumonia caused by *Pasteurella multocida*, and leptospirosis caused by *Leptospira pomona*. In sows, Noromycin 300 LA is indicated as an aid in the control of infectious enteritis (baby pig scours, colibacillosis) in suckling pigs caused by *E. coli*.

Packaging: 100 mL, 250 mL and 500 mL

Administration: IM, IV or SC for specific needs and duration



300 PRO LA _{Rx}

Oxytetracycline 300 mg/mL

300 PRO LA is a ready-to-use injectable, broad-spectrum antibiotic. For beef cattle, non-lactating dairy cattle and calves, including veal calves, 300 PRO LA is indicated for the treatment of pneumonia and shipping fever complex associated with *Pasteurella* spp., and *Histophilus* spp., infectious bovine keratoconjunctivitis (pinkeye) caused by *Moraxella bovis*, footrot and diphtheria caused by *Fusobacterium necrophorum*, bacterial enteritis (scours) caused by *E. coli*, wooden tongue caused by *Actinobacillus lignieresii*, leptospirosis caused by *Leptospira pomona*, and wound infections and acute metritis caused by strains of staphylococcal and streptococcal organisms sensitive to oxytetracycline. Also, 300 PRO LA is indicated for the control of respiratory disease in cattle at high risk of developing BRD associated with *Mannheimia (Pasteurella) haemolytica*. In swine, 300 PRO LA is indicated in the treatment of bacterial enteritis (scours, colibacillosis) caused by *E. coli*, pneumonia caused by *Pasteurella multocida*, and leptospirosis caused by *Leptospira pomona*. In sows, 300 PRO LA is indicated as an aid in the control of infectious enteritis (baby pig scours, colibacillosis) in suckling pigs caused by *E. coli*.

Packaging: 100 mL, 250 mL and 500 mL

Administration: IM, IV or SC for specific needs and duration



Noromectin[®] PLUS

1% w/v Ivermectin and 10% w/v Clorsulon in a Sterile Solution

Noromectin PLUS Injectable ivermectin + clorsulon is a parasiticide for the effective treatment of a wide range of internal and external parasites of beef and non-lactating dairy cattle. Noromectin PLUS offers all the benefits of ivermectin plus control of adult liver flukes. It has FDA approved full "persistent activity" claims and is an economical choice for effective de-worming programs.

Packaging: 50 mL, 100 mL, 250 mL and 500 mL

Administration: SC



Noromectin[®] 1%

1% w/v Ivermectin in a Sterile Solution

Noromectin 1% Injectable ivermectin is a parasiticide for the effective treatment of a wide range of internal and external parasites of beef cattle, non-lactating dairy cattle and swine. It has FDA approved full "persistent activity" claims and is an economical choice for effective de-worming programs.

Packaging: 50 mL, 100 mL, 250 mL and 500 mL

Administration: SC



Norbrook[®]
Pharmaceuticals Worldwide



Noromectin® Pour-On

A Clear Solution Containing Ivermectin 0.5% w/v

Noromectin Pour-On ivermectin offers treatment and control of a wide range of internal and external parasites of beef and non-lactating dairy cattle. Noromectin Pour-On delivers effective control in one convenient low-volume application. It has FDA approved full "persistent activity" claims and is an economical choice for effective lice control with broad-spectrum internal and external parasite control.

Packaging: 1 L, 2.5 L, 5 L and 20 L
Administration: Topical along the mid-line



Oxytet 100

Oxytetracycline 100 mg/mL

Oxytet 100 is effective against a wide range of gram-positive and gram-negative bacteria in beef cattle, beef calves, non-lactating dairy cattle and dairy calves. Oxytet 100 is indicated for the treatment of bacterial pneumonia and shipping fever complex associated with *Pasteurella* spp.; bacterial enteritis (scours) caused by *E. coli*; necrotic pododermatitis (footrot) and calf diphtheria caused by *Fusobacterium necrophorum*; wooden tongue caused by *Actinobacillus lignieresii*; and wound infections, acute metritis and traumatic injury caused by oxytetracycline-susceptible strains of streptococcal and staphylococcal organisms.

Packaging: 500 mL
Administration: IV



Oxytet 200

Oxytetracycline 200 mg/mL

Oxytet 200 is effective against a wide range of gram-positive and gram-negative bacteria in beef cattle, dairy cattle, calves, including pre-ruminating (veal) calves and swine. Oxytet 200 in cattle is indicated for the treatment of pneumonia and shipping fever complex associated with *Pasteurella* spp. and *Haemophilus* spp.; infectious bovine keratoconjunctivitis (pinkeye) caused by *Moraxella bovis*; footrot and diphtheria caused by *Fusobacterium necrophorum*; bacterial enteritis (scours) caused by *E. coli*; wooden tongue caused by *Actinobacillus lignieresii*; leptospirosis caused by *Leptospira pomona*; and wound infections and acute metritis caused by strains of staphylococci and streptococci organisms sensitive to oxytetracycline. In swine, Oxytet 200 is indicated for the treatment of bacterial enteritis (scours, colibacillosis) caused by *E. coli*; pneumonia caused by *Pasteurella multocida*; and leptospirosis caused by *Leptospira pomona*. In sows, Oxytet 200 is indicated as an aid in the treatment of infectious enteritis (baby pig scours, colibacillosis) in suckling pigs caused by *E. coli*.

Packaging: 500 mL
Administration: IM, IV or SC for specific needs and duration



Penicillin Injection

Penicillin G Procaine Injectable Suspension U.S.P.
(300,000 units penicillin G as procaine in a stable suspension)

Penicillin G Procaine is an effective antibacterial agent for use against a variety of pathogenic organisms, chiefly in the gram-positive category, in cattle, sheep, swine and horses.

Packaging: 100 mL and 500 mL
Administration: IM



Flunixin Injection

Flunixin Meglumine Injectable Solution (50 mg/mL) R_x

In cattle, Flunixin is indicated for the control of pyrexia associated with bovine respiratory disease, endotoxemia and acute bovine mastitis and for the control of inflammation in endotoxemia. In horses, Flunixin is indicated for the alleviation of inflammation and pain associated with musculoskeletal disorders especially in acute and chronic stages and for the alleviation of visceral pain associated with colic.

Packaging: 100 mL and 250 mL
Administration: IV and IM in horses and IV in cattle



ALWAYS READ THE ENTIRE PRODUCT INSERT
FOR FULL DISCLOSURE AND WARNINGS PRIOR TO USE.

Contact your distributor for more information or call Norbrook at 913.599.5777.



Noromectin and the Norbrook logo are registered trademarks of Norbrook Laboratories Limited.

Norbrook, Inc.
9733 Loiret Blvd | Lenexa, KS 66219
www.norbrookinc.com